

## Safe and Effective Messaging for Suicide Prevention

### **The Do's—Practices that may be helpful in public awareness campaigns:**

#### **Do emphasize help-seeking and provide information on finding help.**

When recommending mental health treatment, provide concrete steps for finding help. Inform people that help is available through the National Suicide Prevention Lifeline (1-800-273-TALK [8255]) and through established local service providers and crisis centers.

**Do emphasize prevention.** Reinforce the fact that there are preventative actions individuals can take if they are having thoughts of suicide or know others who are or might be. Emphasize that suicides are preventable and should be prevented to the extent possible.<sup>5</sup>

**Do list the warning signs, as well as risk and protective factors of suicide.** Teach people how to tell if they or someone they know may be thinking of harming themselves. Include lists of warning signs, such as those developed through a consensus process led by the [American Association of Suicidology \(AAS\)](#).<sup>6</sup> Messages should also identify protective factors that reduce the likelihood of suicide and risk factors that heighten risk of suicide. Risk and protective factors are listed on pages 35-36 of the [National Strategy for Suicide Prevention](#).

**Do highlight effective treatments for underlying mental health problems.** Over 90 percent of those who die by suicide suffer from a significant psychiatric illness, substance abuse disorder or both at the time of their death.<sup>7-8</sup> The impact of mental illness and substance abuse as risk factors for suicide can be reduced by access to effective treatments and strengthened social support in an understanding community.<sup>9</sup>

### **The Don'ts—Practices that may be problematic in public awareness campaigns:**

**Don't glorify or romanticize suicide or people who have died by suicide.** Vulnerable people, especially young people, may identify with the attention and sympathy garnered by someone who has died by suicide.<sup>10</sup> They should not be held up as role models.

**Don't normalize suicide by presenting it as a common event.** Although significant numbers of people attempt suicide, it is important not to present the data in a way that makes suicide seem common, normal or acceptable. Most people do not seriously consider suicide an option; therefore, suicidal ideation is not normal. Most individuals, and most youth, who seriously consider suicide do not overtly act on those thoughts, but find more constructive ways to resolve them. Presenting suicide as common may unintentionally remove a protective bias against suicide in a community.<sup>11</sup>

**Don't present suicide as an inexplicable act or explain it as a result of stress only.** Presenting suicide as the inexplicable act of an otherwise healthy or high-achieving person may encourage identification with the victim.<sup>12</sup> Additionally, it misses the opportunity to inform audiences of both the complexity and preventability of suicide. The same applies to any explanation of suicide as the understandable response to an individual's stressful situation or to an individual's membership in a group encountering discrimination. Oversimplification of suicide in any of these ways can mislead people to believe that it is a normal response to fairly common life circumstances.<sup>13</sup>

**Don't focus on personal details of people who have died by suicide.** Vulnerable individuals may identify with the personal details of someone who died by suicide, leading them to consider ending their lives in the same way.<sup>14</sup>

**Don't present overly detailed descriptions of suicide victims or methods of suicide.** Research shows that pictures or detailed descriptions of how or where a person died by suicide can be a factor in vulnerable individuals imitating the act. Clinicians believe the danger is even greater if there is a detailed description of the method.<sup>15</sup>